

Resistin and GFR

Kidney International (2006) **70**, 1371. doi:10.1038/sj.ki.5001583

To the Editor: Recently, Axelsson *et al.*¹ showed that glomerular filtration rate was consistently associated with serum resistin levels in patients with chronic kidney disease. It was found that the significant relationship between plasma resistin levels and insulin resistance was lost after adjustment for glomerular filtration rate, leading to the conclusion that renal function is an important factor to take into account when relating insulin sensitivity to inflammatory biomarkers in patients with kidney disease or diabetes mellitus. We have data from a population-based cohort of 64-year-old women, consisting of women with known and new diabetes (diabetes mellitus, $n = 229$), impaired glucose tolerance ($n = 210$), and normal glucose tolerance ($n = 191$).² Diabetes mellitus, impaired glucose tolerance, and normal glucose tolerance were defined according to the World Health Organization classification. The serum levels of resistin were measured with an enzyme-linked immunosorbent assay kit (R&D Systems Europe, Abingdon, UK). The characteristics of the subjects are presented in Table 1. Serum creatinine did not differ

Table 1 | Characteristics of the subjects (geometric means \pm s.d.) ($n=630$)

	DM ($n=229$)	IGT ($n=210$)	NGT ($n=191$)
Serum resistin (ng/ml)	10.8 ± 3.6	10.2 ± 3.7	$9.9 \pm 3.2^{**}$
Serum creatinine ($\mu\text{mol/l}$)	89 ± 14	89 ± 11	89 ± 10
BMI (kg/m^2)	28.8 ± 4.9	27.1 ± 5.0	$25.8 \pm 3.5^*$
Blood glucose (mmol/l)	7.1 ± 2.8	5.0 ± 0.6	$4.7 \pm 0.5^*$
Serum HDL cholesterol (mmol/l)	1.45 ± 0.43	1.56 ± 0.42	$1.74 \pm 0.42^*$
White blood cell count ($\times 10^9/\text{l}$)	6.6 ± 1.8	6.1 ± 1.8	$5.6 \pm 1.5^*$
C-reactive protein (mg/l)	1.9 ± 10.3	1.3 ± 3.7	$1.3 \pm 2.5^*$

Abbreviations: BMI, body mass index; DM, diabetes mellitus; HDL, high-density lipoprotein; IGT, impaired glucose tolerance; NGT, normal glucose tolerance; s.d., standard deviation.

* $P < 0.001$; ** $P = 0.006$.

between the groups and correlated positively to serum resistin (Spearman's correlation coefficients 0.36 ($P < 0.01$), 0.28 ($P < 0.01$), and 0.15 ($P < 0.05$) for the diabetes mellitus, impaired glucose tolerance, and normal glucose tolerance groups, respectively). In a multivariate analysis, serum creatinine remained associated with resistin after adjustment for C-reactive protein, white blood cell count, high-density lipoprotein cholesterol, body mass index, glucose tolerance group, and plasma insulin (partial correlation coefficient 0.33, $P < 0.001$). These data indicate that the association between glomerular filtration rate and serum resistin also represents a physiological mechanism that not necessarily is linked to kidney disease.

1. Axelsson J, Bergsten A, Qureshi AR *et al.* Elevated resistin levels in chronic kidney disease are associated with decreased glomerular filtration rate and inflammation, but not with insulin resistance. *Kidney Int* 2006; **69**: 596–604.
2. Brohall G, Behre CJ, Hulthe J *et al.* Prevalence of diabetes and impaired glucose tolerance in 64-year-old Swedish women. *Diabetes Care* 2006; **29**: 363–367.

B Fagerberg¹, C Fagerlund¹ and J Hulthe¹

¹Wallenberg Laboratory for Cardiovascular Research, Sahlgrenska University Hospital, Göteborg, Sweden

Correspondence: B Fagerberg, Wallenberg Laboratory for Cardiovascular Research, Sahlgrenska University Hospital, Göteborg SE 41345, Sweden.

E-mail: bjorn.fagerberg@wlab.gu.se

Is there a link between inflammation, plasma resistin levels, and protein malnutrition in hemodialysis patients?

Kidney International (2006) **70**, 1371–1372. doi:10.1038/sj.ki.5001584

To the Editor: Similarly to the article by Axelsson *et al.*¹ entitled 'Elevated resistin levels in chronic kidney disease are associated with decreased glomerular filtration rate and inflammation, but not with insulin resistance' we have recently observed in 33 non-diabetic hemodialysis patients that elevated resistin plasma levels are not associated with insulin resistance. In addition, we detected for the first time in human subjects that resistin levels are inversely correlated with the protein catabolic rate ($r = -0.416$, $P = 0.025$), which is known to parallel the dietary protein uptake (Figure 1).² Axelsson *et al.*¹ have not reported on this important aspect in their manuscript despite the high prevalence of protein and energy malnutrition in hemodialysis patients and in patients with chronic kidney disease.

The finding by Axelsson *et al.*¹ that plasma resistin levels are strongly correlated with inflammatory markers and




Figure 1 | Hemodialysis patients ($n = 33$) were divided into three groups of 11 patients each according to the protein catabolic rate. The patients with the lowest protein catabolic rate exhibited the highest resistin levels. These intergroup differences are statistically significant ($P < 0.05$).

independent from glomerular filtration rates is of considerable interest, as elevated resistin levels in various other disorders have been associated with both inflammation as well as malnutrition.^{3–5}

Taking into account that inflammation has been closely related to malnutrition and in particular to protein–energy malnutrition,⁶ one may assume that there is a relationship between inflammation, plasma resistin levels, and protein malnutrition in hemodialysis patients and in patients with chronic kidney disease that is yet to be clarified.

1. Axelsson J, Bergsten A, Qureshi AR *et al.* Elevated resistin levels in chronic kidney disease are associated with decreased glomerular filtration rate and inflammation, but not with insulin resistance. *Kidney Int* 2006; **69**: 596–604.
2. Filippidis G, Liakopoulos V, Mertens PR *et al.* Resistin serum levels are increased but not correlated with insulin resistance in chronic hemodialysis patients. *Blood Purif* 2005; **23**: 421–428.
3. Karmiris K, Koutroubakis IE, Xidakis C *et al.* Circulating levels of leptin, adiponectin, resistin, and ghrelin in inflammatory bowel disease. *Inflamm Bowel Dis* 2006; **12**: 100–105.
4. Housova J, Wilczek H, Haluzik MM *et al.* Adipocyte-derived hormones in heroin addicts: the influence of methadone maintenance treatment. *Physiol Res* 2005; **54**: 73–78.
5. Bahr MJ, Ockenga J, Boker KH *et al.* Elevated resistin levels in cirrhosis are associated with the proinflammatory state and altered hepatic glucose metabolism, but not with insulin resistance. *Am J Physiol Endocrinol Metab* 2006 (in press, doi:10.1152/ajpendo.00291.2005).
6. Sauerwein RW, Mulder JA, Mulder L *et al.* Inflammatory mediators in children with protein–energy malnutrition. *Am J Clin Nutr* 1997; **65**: 1534–1539.

V Liakopoulos¹, PR Mertens², T Eleftheriadis¹, G Koukoulis³ and I Stefanidis¹

¹Department of Nephrology, University of Thessalia School of Medicine, Larissa, Greece; ²Department of Nephrology and Immunology, University Hospital Aachen, Aachen, Germany and ³Department of Endocrinology, University of Thessalia School of Medicine, Larissa, Greece

Correspondence: I Stefanidis, Department of Nephrology, University of Thessalia School of Medicine, Mezourlo Hill, Larissa 41110, Greece.

E-mail: stefanid@med.uth.gr

Response to ‘Resistin letters’

Kidney International (2006) **70**, 1372. doi:10.1038/sj.ki.5001602

We thank Fagerberg *et al.* for their contribution to the growing body of evidence refuting a causative relationship between the protein resistin and insulin resistance in man.¹ Indeed, resistin was first described as found in inflammatory zone (FIZZ)-1, indicating its structural similarity to several other proinflammatory cytokines.² Indeed, resistin is expressed mainly in immunocompetent cells such as leukocytes, and not in adipocytes as had previously been thought.^{3,4}

Given the fact that even a mild renal function impairment is associated with a chronic low-grade inflammation⁵ and a markedly increased risk of cardiovascular death, it is not surprising that the patients in the study of Fagerberg *et al.* displayed increased resistin levels that were also correlated with serum creatinine, C-reactive protein, and leukocyte counts.

Taken together, we believe that this study further underscores the relationship between decreasing renal function and increased risks of inflammation and cardiovascular disease. Resistin is one part of this complex, where we are still looking for a unifying mechanism and an effective therapy.

1. Arner P. Resistin: yet another adipokine tells us that men are not mice. *Diabetologia* 2005; **48**: 2203–2205.
2. Holcomb IN, Kabakoff RC, Chan B *et al.* FIZZ1, a novel cysteine-rich secreted protein associated with pulmonary inflammation, defines a new gene family. *EMBO J* 2000; **19**: 4046–4055.
3. Patel L, Buckels AC, Kinghorn IJ *et al.* Resistin is expressed in human macrophages and directly regulated by PPAR gamma activators. *Biochem Biophys Res Commun* 2003; **300**: 472–476.
4. Sopasakis VR, Nagaev I, Smith U. Cytokine release from adipose tissue of nonobese individuals. *Int J Obes (London)* 2005; **29**: 1144–1147.
5. Fried L, Solomon C, Shlipak M *et al.* Inflammatory and prothrombotic markers and the progression of renal disease in elderly individuals. *J Am Soc Nephrol* 2004; **15**: 3184–3191.

J Axelsson¹, B Lindholm¹ and P Stenvinkel¹

¹Department of Renal Medicine, Karolinska Institute, K56 Karolinska University Hospital at Huddinge, Stockholm, Sweden

Correspondence: J Axelsson, Department of Renal Medicine, Karolinska Institute, K56 Karolinska University Hospital at Huddinge, Stockholm 141 86, Sweden. E-mail: jonas.axelsson@ki.se

Enlightenment on liver lanthanum exposure

Kidney International (2006) **70**, 1372–1373. doi:10.1038/sj.ki.5001636

To the Editor: Tissue lanthanum concentrations in the paper by Slatopolsky *et al.*¹ are similar to our own and others’ findings,² but have been interpreted very differently. Comparisons with ‘untreated’ controls are used to highlight ‘marked’ and ‘striking’ increases in deposition in nephrectomized rats, including a 98-fold increase in liver concentration. When the appropriate comparator is used (lanthanum-treated rats with normal renal function), there is no statistical increase in bone and kidney concentrations in the uremic animals, and only two- to three-fold higher liver concentrations after 15 weeks of treatment. This is entirely consistent with the absence of any effect of renal failure on systemic lanthanum exposure³ and a small (two- to three-fold) increase in first-pass liver exposure, as reported recently for this uremic model.² Although tissue concentrations increased over time (termed ‘progressive accumulation’), their study was too short to fully reflect the tissue kinetics of lanthanum, and longer-term studies suggest achievement of steady-state conditions (Figure 1). Although a maximal dose of lanthanum (3% (w/w) of diet, 15–22 times the human dose of 3 g/day)¹ was administered for as long as practicable in a nephrectomized model (limited by death owing to renal failure), lanthanum concentrations were still only at trace levels (<3 p.p.m.). As the natural liver concentrations of other lysosomally transported metals such as iron and copper are up to 2007 and 55 p.p.m., respectively,⁴ there would